

Camille M Clinton, MD
ProOrtho
12911 120th Ave NE, Suite H-210
Kirkland, WA 98034

Shoulder Anterior Capsulorrhaphy / Bankart Repair

0-4 weeks post op

- Arm in sling/immobilizer
- Modalities as needed
- PROM with ATC or PT no shoulder pulleys
 **Flexion to 90°, Abduction to 90°, Internal Rotation to 30° in scapular plane,
 External Rotation to neutral And Extension to neutral**
- Isometric strengthening and ROM of the hand, wrist, and elbow

4 weeks post op → begin formal physical therapy

- D/C sling/immobilizer
- No IR or ER resistance exercises
- Periscapular isometrics
- PROM with shoulder pulleys
 **Flexion to 130°, Extension to 30°, Abduction to 130°, Internal
 Rotation to 90° in plane of scapula, And External Rotation 30° in
 plane of scapula**
- Progress from AAROM to AROM:
 - Quality movement only, **avoid forcing** active motion with substitution patterns.
 - Remember the effects of gravity on the limb, do gravity eliminated motions first ie. Supine flexion.
- Deltoid isometrics.
- Lightly resisted **elbow** flexion

6 weeks post op

- Advance ROM to full as tolerated, do not crank on ER but progress it as tolerated,
- Begin standing isotonic RC exercises with arm in neutral, below 90°
- Continue periscapular isometrics
- Joint mobilization (posterior glides)

Shoulder Anterior Capsulorrhaphy / Bankart Repair (page 2)

10 weeks post op

- Full A/PROM in all direction with normal ratio of movement between GH joint and SC joint
- Advance Rotator cuff strengthening to 8-10lbs on all motions
- Advance proprioception ex's as tolerated

12 weeks post op

- Add Eccentric ex's to posterior cuff with T-band in standing simulating follow through in throwing or
 sidelying ER

Camille M Clinton, MD
ProOrtho
12911 120th Ave NE, Suite H-210
Kirkland, WA 98034

- Full ROM isokinetics
- May begin conventional weight lifting using machines and progressing to free weights if desired as tolerated

14 weeks post op

- Begin light tennis ball tossing at 20-30ft. max at 60% velocity, work on mechanics of wind up, early cocking phase, late cocking phase, acceleration, and follow through

16-24 weeks post op

- Throwers must pass strength test to begin throwing
- Retest monthly till strength test passed
- Continue with maintenance strengthening

Return to Sport/Activity

- When cleared by physician
- Pass strength test
- Throwers complete throwing program
- No pain with all desired level of activities
- 4-6 months